

# Omega photoproduction beyond the resonance region at small $u$

A. Sibirtsev<sup>1</sup>, K. Tsushima<sup>1,2,3</sup> and S. Krewald<sup>1,2</sup>

<sup>1</sup>*Institut für Kernphysik,*

*Forschungszentrum Jülich, D-52425 Jülich*

<sup>2</sup>*Special Research Center for the Subatomic Structure of Matter (CSSM) and Department of Physics and Mathematical Physics, University of Adelaide, SA 5005, Australia*

<sup>3</sup>*Department of Physics and Astronomy, University of Georgia, Athens, GA 30602 USA*

Within the meson exchange model we study  $\omega$  meson photoproduction at energies above the  $s$  channel resonance region. Different model prescriptions for the  $\omega NN$  vertex function are investigated imposing gauge invariance as well as crossing symmetry. The calculations reproduce the energy dependence of the differential  $\omega$  photoproduction cross sections at moderate  $|u|$  for  $E_\gamma \leq 4.7$  GeV, which previously were discussed as an indication of the hard interaction between the photon and quarks of the nucleon.

PACS numbers: PACS: 13.60.Le; 13.88.+e; 14.20.Gk; 25.20.Lj

## I. INTRODUCTION

At energies beyond the  $s$  channel resonance region the vector meson photoproduction off the nucleon is traditionally discussed in terms of exchanges in the  $t$  and  $u$  channels.

At forward angles the vector meson photoproduction is dominated by the exchanges in the  $t$  channel. A long time ago, Berman and Drell [1] proposed that at low energies and small  $t$  meson photoproduction can be well understood in terms of the exchanges of light mesons.

Most recently, Donnachie and Landshoff [2, 3] and Laget [4] illustrated that at high energies the photoproduction can be well described by soft pomeron and meson Regge trajectories.

Furthermore, the high energy data [5] on  $\rho$  photoproduction at  $|t|$  above  $\simeq 0.4$  GeV<sup>2</sup> require some additional contribution, which might stem from the hard pomeron exchange [6]. The data [7] on  $J/\Psi$  meson photoproduction can be reproduced well by an additional introduction of the hard pomeron exchange in the  $t$  channel [6, 8, 9] even at small  $|t|$ .

At backward angles, i.e. where  $|t|$  becomes large and  $|u|$  still remains small, the photoproduction is dominated by exchanges in the  $u$  channel. At low energies beyond the resonance region, the production amplitude is given by the nucleon exchange. At high energies it is due to the contribution from the nucleon Regge trajectory.

Meson photoproduction at the kinematical region where both  $t$  and  $u$  are large can be dominated by hard interactions between the photon and the quarks of the nucleon. It is believed that this region is beyond the applicability of the standard boson exchange and Regge models. Moreover, a typical signature [11] of hard processes is a strong  $\propto s^{-8}$  dependence of the differential photoproduction cross section.

Experimental results [10] on backward  $\omega$  meson photoproduction at photon energies  $E_\gamma$  between 2.8 and

4.8 GeV collected at Daresbury Laboratory indicate a prominent structure shown in Fig. 1. The differential  $d\sigma/du$  cross section has a dip around  $u \simeq 0.15$  GeV<sup>2</sup>, which becomes more pronounced with increasing the photon energy.

It was proposed [10] to parameterize the  $\gamma p \rightarrow \omega p$  differential cross section as a sum of contributions from the Regge trajectory exchange and hard mechanism. Indeed the energy dependence of the  $\omega$  meson photoproduction cross section  $d\sigma/du$  at different values of  $u$  was fitted well by

$$\frac{d\sigma}{du} = |a(u)s^{\alpha(u)-1} + b(u)\exp^{i\phi(u)}s^{-n/2}|^2, \quad (1)$$

with fixed power  $n=8$  and parameters  $a$ ,  $b$  and  $\phi$  adjusted to the data [10]. It was found that  $\alpha(u)$  corresponds to the nucleon trajectory, that the parameter  $b$  does not depend on  $u$  and the phase  $\phi=90^\circ$  indicating that the Regge exchange and hard mechanism do not interfere.

Finally, the data [10] on backward  $\omega$  photoproduction off a nucleon at  $E_\gamma$  between 2.8 and 4.8 GeV were interpreted as due to the dominant contribution from the nucleon Regge trajectory exchange at  $|u| \leq 0.2$  GeV<sup>2</sup> and an additional contribution from the hard interaction between the photon and the quarks of the target nucleon.

On the other hand, the data [12] on backward  $\rho$  and  $f$  meson photoproduction collected at the same spectrometer and for the same range of photon energies do not indicate a dip around  $u \simeq -0.15$  GeV<sup>2</sup> as well as a strong  $s^{-8}$  dependence.

Furthermore high precision measurements [13] at SLAC of the backward photoproduction of charged pions off a nucleon at photon energies between 4.1 and 14.8 GeV and  $-1.8 \leq u \leq 0.05$  GeV<sup>2</sup> also do not indicate a dip at  $u \simeq -0.15$  GeV<sup>2</sup>. SLAC data [14] on backward  $\pi^0$  photoproduction at photon energies  $E_\gamma=6, 8, 12$  and 18 GeV show a  $s^{-3}$  dependence with no sign of a dip at  $u$  between -1 and 0 GeV<sup>2</sup>.

Therefore, among available data on meson photoproduction at large angles only the  $\gamma p \rightarrow \omega p$  reaction shows a very specific signature which might be interpreted as an indication of hard interactions between the photon and the nucleon. However it is quite difficult to motivate that the  $\omega$  photoproduction can be sensitive to hard processes whereas photoproduction of  $\pi$ ,  $\rho$  and  $f$  mesons are not, since hard interactions are driven by the partonic structure of the hadrons.




FIG. 1: The data [10] on the differential  $\gamma p \rightarrow \omega p$  cross section as function of  $u$  for different photon energies  $E_\gamma$ . The lines show our meson exchange calculations with different prescriptions for the  $\omega NN$  vertex functions given by models A (dashed), B (dotted), C (dot-dashed), and D (solid).

Here we investigate whether the data [10] on backward  $\omega$  meson photoproduction might be understood in terms of a standard hadronic approach as a meson exchange model. At photon energies above the  $s$  resonance region we adopt the  $\pi$ ,  $\eta$  and  $\sigma$  exchanges in the  $t$  channel as well as the nucleon exchange in the  $s$  and  $u$  channels. Within meson exchange models, the contribution from the nucleon exchange dominates at large angles.

It is important to note that within the meson exchange model, the difference between the data on backward  $\omega$  and  $\pi$ ,  $\rho$  and  $f$  meson photoproduction can be reasonably motivated, since the couplings and the form factor functions in the meson-nucleon-nucleon vertices are not

necessarily identical for the various mesons. Moreover, theoretical analysis [15, 16] of the nucleon-nucleon interaction really result in different vertex functions for different mesons.

## II. THE MODEL

The vector meson photoproduction at small four momentum  $t$  transfer has been studied theoretically for a long time [17, 18, 19]. Considering the contributions from all possible intermediate states that are lighter than the produced vector mesons, Joos and Kramer [19] formulated the  $\pi$ ,  $\eta$  and  $\sigma$  exchange model.

The coupling constants for the  $\pi$ ,  $\eta$  and  $\sigma$  exchanges were fixed either from the analysis of nucleon-nucleon scattering or from the relevant partial decay widths of the vector meson to the photon and the pseudoscalar meson. The  $\omega\gamma\sigma$  coupling was not specified explicitly. Form factors in the interaction vertices were not taken into account [20, 21].

However, the model in its original formulation [1, 17, 18, 19] was not able to reproduce the data on  $\omega$  photoproduction at energies  $1.8 \leq E_\gamma \leq 5.8$  GeV collected later [22, 23, 24, 25]. We notice that the  $\omega$  photoproduction data [1, 17, 18, 19] at low  $|t|$  can be well described by the model of Ref. [19] when introducing the relevant form factors at the interaction vertices.

Following the model of Refs. [1, 17, 18, 19] and evaluating the interaction vertices by vector meson dominance Friman and Soyeur [27] reproduced the data [26] on total  $\gamma p \rightarrow \omega p$  cross section at photon energies below 2 GeV and differential  $d\sigma/dt$  omega photoproduction cross sections [22] at  $|t| \leq 0.5$  GeV<sup>2</sup> for  $1.4 \leq E_\gamma \leq 1.8$  GeV. In this model [27] the photon couples to the  $\rho$  and the  $\omega$  and the  $\rho N \rightarrow \omega N$  transition is given by  $\pi$  meson exchange, while the  $\omega N \rightarrow \omega N$  scattering was described by  $\sigma$  exchange. Furthermore, the interaction vertices were dressed by form factors and unknown parameters for the  $\sigma$  meson exchange were adjusted to the data [22].

Obviously, since both models [19, 27] discussed above account only for the  $t$ -channel contribution, they are not able to reproduce the data [10] collected at small  $|u|$ .

Very recently Oh, Titov and T.S.H. Lee [28] investigated  $\omega$  photoproduction in the resonance region by considering the  $\pi$ ,  $\eta$  and pomeron exchanges in the  $t$  channel and the excitation of the nucleon and baryonic resonances in the  $s$  and  $u$  channels. The form factors and coupling constants for  $\pi$ ,  $\eta$  and nucleon exchanges were adopted from the literature. The parameters for the pomeron exchange were fixed [2, 29, 30] at high energies. Furthermore, the resonance parameters were fitted to the  $\gamma p \rightarrow \omega p$  differential cross sections [31] collected at photon energies  $1.23 \leq E_\gamma \leq 1.92$  GeV. The calculations [28] also well reproduce the data [32] at small  $|t|$  for  $E_\gamma = 2.8$  and 4.7 GeV. However, the model could not reproduce the  $\omega$  photoproduction data [10] at small  $|u|$ .

Here we consider  $\omega$  meson photoproduction due to the

pseudoscalar  $\pi$  and  $\eta$  and scalar  $\sigma$  meson exchanges and the direct and crossed nucleon terms. The relevant diagrams are shown in the Fig. 2. Furthermore, the squared invariant collision energy  $s$  and the squared  $t$  and  $u$  four momenta transfers are given by

$$s = (p + k)^2 = (p' + q)^2, \quad (2)$$

$$t = (q - k)^2 = (p' - p)^2, \quad (3)$$

$$u = (q - p)^2 = (p' - k)^2, \quad (4)$$

where  $k$ ,  $q$ ,  $p$ , and  $p'$  are the four momenta of the photon,  $\omega$  meson, initial nucleon and final nucleon, respectively.




FIG. 2:  $\omega$  meson photoproduction mechanism in  $t$  (a),  $u$  (b) and  $s$  (c) channels.

The effective Lagrangian densities used for the evaluation of the  $\pi$  and  $\eta$  meson exchange amplitudes are given as

$$\begin{aligned} \mathcal{L}_{\omega\gamma\varphi} &= \frac{eg_{\omega\gamma\varphi}}{m_\omega} \epsilon^{\mu\nu\alpha\beta} \partial_\mu \omega_\nu \partial_\alpha A_\beta \varphi, \\ \mathcal{L}_{\varphi NN} &= -ig_{\varphi NN} \bar{N} \gamma_5 N \varphi, \end{aligned} \quad (5)$$

where  $\varphi$  denotes the  $\pi^0$  or the  $\eta$  meson, while  $A_\beta$  is the photon field. Furthermore we adopt the  $\pi NN$  and  $\eta NN$  coupling constants as  $g_{\pi NN}=13.26$  and  $g_{\eta NN}=3.53$ , respectively. The  $g_{\omega\gamma\pi}=1.82$  and  $g_{\omega\gamma\eta}=0.42$  coupling constants were evaluated from the partial  $\omega \rightarrow \gamma\pi$  and  $\omega \rightarrow \gamma\eta$  decay widths [33], respectively.

Finally, the  $\gamma p \rightarrow \omega p$  reaction amplitude due to the  $\pi$  and  $\eta$  meson exchange is given as

$$\begin{aligned} \mathcal{M}_{\pi,\eta} &= \sum_{\varphi=\pi,\eta} \frac{-ieg_{\omega\gamma\varphi}g_{\varphi NN}}{m_\omega} \frac{1}{t-m_\varphi^2} \\ &\times \epsilon^{\mu\nu\alpha\beta} q_\mu \epsilon_\nu^*(q) k_\alpha \epsilon_\beta(k) \bar{u}(p') \gamma_5 u(p). \end{aligned} \quad (6)$$

Moreover, at each interaction vertex we introduce the phenomenological form factor

$$F(t) = \frac{\Lambda^2 - m_{ex}^2}{\Lambda^2 - t}, \quad (7)$$

where  $m_{ex}$  is the mass of the exchanged meson and the cutoff parameters in the  $\pi NN$  and  $\eta NN$  vertices were fixed as  $\Lambda_{\pi NN}=0.7$  GeV and  $\Lambda_{\eta NN}=1$  GeV. The cutoff parameters of the form factors in the  $\omega\gamma\pi$  and  $\omega\gamma\eta$  vertices were fitted to the data as will be specified later.

The  $\sigma$  meson exchange amplitude was calculated from the following effective Lagrangians:

$$\begin{aligned} \mathcal{L}_{\omega\gamma\sigma} &= em_\omega g_{\omega\gamma\sigma} \omega_\beta A^\beta \sigma \\ \mathcal{L}_{\sigma NN} &= g_{\sigma NN} \bar{N} N \sigma, \end{aligned} \quad (8)$$

where the  $\sigma NN$  coupling constant was taken as  $g_{\sigma NN}=8$ , while the  $\omega\gamma\sigma$  coupling constant was fitted to the data.

The photoproduction amplitude due to the  $\sigma$  meson exchange is then naively given as

$$\mathcal{M}_\sigma = \frac{em_\omega g_{\omega\gamma\sigma} g_{\sigma NN}}{t - m_\sigma^2} \epsilon_\mu^*(q) \epsilon^\mu(k) \bar{u}(p') u(p). \quad (9)$$

Again we introduce monopole form factor of Eq. (7) in the interaction vertices with  $\Lambda_{\sigma NN}=1$  GeV and adjust the cutoff  $\Lambda_{\omega\gamma\sigma}$  to the data. Furthermore, the gauge invariant  $\sigma$  exchange amplitudes is now given as

$$\begin{aligned} \mathcal{M}_\sigma &= \frac{em_\omega g_{\omega\gamma\sigma} g_{\sigma NN}}{t - m_\sigma^2} \\ &\times \epsilon_\mu^*(q) \left( g^{\mu\nu} - \frac{k^\mu q^\nu}{q \cdot k} \right) \epsilon_\nu(k) \bar{u}(p') u(p). \end{aligned} \quad (10)$$

The direct and crossed nucleon exchange amplitudes due to the diagrams depicted in Fig. 2c) and Fig. 2b), respectively, were evaluated from the following interaction Lagrangians:

$$\begin{aligned} \mathcal{L}_{\gamma NN} &= -e\bar{N} \left( \gamma_\mu \frac{1+\tau_3}{2} A^\mu - \frac{\kappa_p}{2M_N} \sigma^{\mu\nu} \partial_\nu A_\mu \right) N, \\ \mathcal{L}_{\omega NN} &= -g_{\omega NN} \bar{N} \left( \gamma_\mu \omega^\mu - \frac{\kappa_\omega}{2M_N} \sigma^{\mu\nu} \partial_\nu \omega_\mu \right) N, \end{aligned} \quad (11)$$

where the anomalous magnetic moment of the proton was taken as  $\kappa_p=1.79$ , while the  $\omega NN$  coupling constant was fitted to the data. Moreover, in the following calculations we adopt  $\kappa_\omega=0$ .

Ignoring the form factors for the  $\gamma p \rightarrow \omega p$  reaction, the reaction amplitudes due to the nucleon exchange in the  $s$  and  $u$  channels are given as

$$\begin{aligned} \mathcal{M}_N &= eg_{\omega NN} \bar{u}(p') \epsilon_\mu^*(q) \gamma^\mu \frac{\not{p} + \not{k} + m_N}{s - m_N^2} \\ &\times [\gamma_\nu + \frac{i\kappa_p}{2m_N} \sigma_{\nu\alpha} k^\alpha] \epsilon^\nu(k) u(p), \end{aligned} \quad (12)$$

$$\begin{aligned} \tilde{\mathcal{M}}_N &= eg_{\omega NN} \bar{u}(p') \epsilon_\mu^*(q) [\gamma_\nu + \frac{i\kappa_p}{2m_N} \sigma_{\nu\alpha} k^\alpha] \\ &\times \frac{\not{p} - \not{q} + m_N}{u - m_N^2} \gamma^\mu \epsilon^\nu(k) u(p). \end{aligned} \quad (13)$$

The form factor in the  $\omega NN$  vertex will be specified in the next section.

Finally, the model parameters which we fixed prior to the global fitting to the data are listed in Table I. However let us to remind that for each meson exchange diagram shown in the Fig.2a) the form factors in the  $\omega\gamma\pi$ ,  $\omega\gamma\eta$  and  $\omega\gamma\sigma$  vertices were adjusted to the data, while those in the  $\pi NN$ ,  $\eta NN$  and  $\sigma NN$  vertices, respectively, were fixed. Since the  $\mathcal{M}_\pi$ ,  $\mathcal{M}_\eta$  and  $\mathcal{M}_\sigma$  amplitudes contain the product of the form factors in both vertices, the model could not be used for a unique determination of the relevant vertex function parameters.

TABLE I: Coupling constants  $g$  and cut-off parameters  $\Lambda$  fixed in our calculations. The cut-off parameters are given for the form factor of Eq. (7).

| vertex             | $g$   | $\Lambda$ (GeV) |
|--------------------|-------|-----------------|
| $\pi NN$           | 13.26 | 0.7             |
| $\eta NN$          | 3.53  | 1.0             |
| $\sigma NN$        | 8.0   | 1.0             |
| $\omega\gamma\pi$  | 1.82  | —               |
| $\omega\gamma\eta$ | 0.42  | —               |

### III. THE $\omega NN$ VERTEX FUNCTION

A naive evaluation of the amplitude and/or the introduction of phenomenological form factors in the photoproduction reactions at the level of Born amplitudes violates gauge invariance. There are a few common recipes [34, 35, 36, 37, 38] proposed to restore gauge invariance. The dominant contribution at small  $u$  comes from the nucleon exchange current. Therefore, the data on  $\omega$  photoproduction at small  $u$  should be sensitive to the prescription of the  $\omega NN$  vertex function as well as to the methods for restoration of gauge invariance of the nucleon exchange amplitude.

In order to understand whether it is necessary to use form factors in the  $\omega NN$  vertices at  $s$  and  $u$  channels of the diagrams (b) and (c) in Fig.2 we fit the data without  $\omega NN$  form factors and vary the  $g_{\omega NN}$  and  $g_{\omega\gamma\sigma}$  coupling constants and cut-off parameters at the  $\omega\gamma\pi$ ,  $\omega\gamma\eta$  and  $\omega\gamma\sigma$  vertices. We denote these calculations as Model A.

Furthermore, we use simultaneously the  $\gamma p \rightarrow \omega p$  data on the  $d\sigma/du$  [10] and the  $d\sigma/dt$  differential cross sections available at photon energies  $2.8 \leq E_\gamma \leq 4.7$  GeV. The global fit was performed with Minuit [39] and includes 120 experimental points.

The results obtained with model A are shown by the dashed lines in Fig. 1 and Fig. 3. The relevant sets of model parameters and the total  $\chi^2$  are listed in Table II. The calculations describe the data at small  $|u|$  at  $E_\gamma=4.7$  GeV well. However, model A could not reproduce the dependence of the  $\omega$  photoproduction differential cross section  $d\sigma/du$  on the photon energy. Moreover, the fit results in large  $\omega\gamma\sigma$  and very small  $\omega NN$  coupling constants. Therefore we consider the parameters

TABLE II: Model parameters fitted to the data [10]. Model A is the calculations without form factor at the  $\omega NN$  vertex. Model B stands for the calculations with form factor of Eq. (14) at the  $\omega NN$  vertex. Model C denotes the calculations with the form factor of Eq. (17) at the  $\omega NN$  vertex. Model D denotes the calculations with the form factor of Eq. (18). The cut-off parameters are given in GeV. The  $\chi^2$  is total chi-square.

| parameter                      | model A         | model B         | model C         | model D         |
|--------------------------------|-----------------|-----------------|-----------------|-----------------|
| $\Lambda_{\omega\gamma\pi}$    | $0.10 \pm 0.07$ | $0.10 \pm 0.04$ | $0.40 \pm 0.01$ | $0.60 \pm 0.03$ |
| $\Lambda_{\omega\gamma\eta}$   | $0.1 \pm 0.3$   | $0.1 \pm 0.6$   | $0.4 \pm 0.1$   | $1.4 \pm 0.9$   |
| $g_{\omega\gamma\sigma}$       | $7.2 \pm 0.2$   | $10 \pm 1$      | $1.4 \pm 0.2$   | $1.11 \pm 0.01$ |
| $\Lambda_{\omega\gamma\sigma}$ | $0.59 \pm 0.01$ | $0.52 \pm 0.01$ | $0.83 \pm 0.07$ | $1.0 \pm 0.1$   |
| $g_{\omega NN}$                | $0.22 \pm 0.01$ | $0.26 \pm 0.02$ | $5.8 \pm 0.5$   | $1.7 \pm 0.1$   |
| $\Lambda_{\omega NN}$          | —               | $2.21 \pm 0.01$ | $0.30 \pm 0.01$ | $1.85 \pm 0.03$ |
| $\tilde{\Lambda}_{\omega NN}$  | —               | —               | $1.23 \pm 0.03$ | —               |
| $\chi^2$                       | 1796            | 1799            | 614             | 598             |

of model A as unreasonable. Finally we conclude that the calculations without a form factor at the  $\omega NN$  vertex cannot not reproduce the complete set of data on the  $\gamma p \rightarrow \omega p$  reaction.

Now we have followed Refs. [28, 37] and included a form factor at the  $\omega NN$  vertex in the  $u$  and  $s$  channels

$$F(r) = \frac{\Lambda^4}{\Lambda^4 + (r - m_N^2)^2}, \quad (14)$$

where  $r=s$  or  $u$ . The cut-off parameter  $\Lambda$  was fitted to the data.

Furthermore, the gauge invariant nucleon exchange amplitudes are now given as

$$\begin{aligned} \mathcal{M}_N &= eg_{\omega NN} \bar{u}(p') \varepsilon_\mu^*(q) [\gamma^\mu - k^\mu \frac{\not{q}}{q \cdot k}] \frac{\not{p} + \not{k} + m_N}{s - m_N^2} \\ &\times F(s) [\gamma_\nu + \frac{\kappa_p}{2m_N} \gamma_\nu \not{k} - \frac{q_\nu}{q \cdot k} \not{k}] \varepsilon^\nu(k) u(p), \end{aligned} \quad (15)$$

$$\begin{aligned} \tilde{\mathcal{M}}_N &= eg_{\omega NN} \bar{u}(p') \varepsilon_\mu^*(q) [\gamma_\nu + \frac{\kappa_p}{2m_N} \gamma_\nu \not{k} - \frac{q_\nu}{q \cdot k} \not{k}] \\ &\times F(u) \frac{\not{p} - \not{q} + m_N}{u - m_N^2} [\gamma^\mu - k^\mu \frac{\not{q}}{q \cdot k}] \varepsilon^\nu(k) u(p). \end{aligned} \quad (16)$$

We denote the calculations with the form factor of Eq. (14) as model B and show the results by the dotted lines in Fig. 1 and Fig. 3. The parameters of model B are given in Table II.

The calculations by model B again could not reproduce the dependence of the  $\gamma p \rightarrow \omega p$  differential cross section on the photon energy. The data on  $d\sigma/dt$  and  $d\sigma/du$  at  $E_\gamma=4.7$  GeV are well described, however. We also notice that our results shown in Fig. 1 are different from those reported in Ref. [28]. We attribute this discrepancy to the fact that in our study the complete set of available  $\omega$  photoproduction data on both  $d\sigma/dt$  and  $d\sigma/du$  is included for the whole energy range  $2.8 \leq E_\gamma \leq 4.7$  GeV,

while Oh, Titov and T.S.H. Lee [28] adjusted the model parameters to the  $d\sigma/dt$  data only.

In order to solve the so-called A2-problem of pion photoproduction, Davidson and Workman [38] recently suggested a new kind of form factor motivated by gauge invariance and crossing symmetry which reads as follows:

$$F(u, s) = F(u, \Lambda) + F(s, \tilde{\Lambda}) - F(u, \Lambda)F(s, \tilde{\Lambda}), \quad (17)$$

with the form factors taken in the form of Eq. (14). The cut-off parameters  $\Lambda$  and  $\tilde{\Lambda}$  should be identical in order to satisfy crossing symmetry. The calculations performed with this form factor are denoted as model C. The results are displayed in Fig. 1 and Fig. 3 by the dashed-dotted lines. The relevant parameters are listed in Table II. It turned out that the crossing symmetry had to be broken, if a good fit to the data was required. The question arises whether the idea of Davidson and Workman should be abandoned.

We found that a mild modification of the ansatz of Davidson and Workman allows to maintain the crossing symmetry. We investigate the ansatz:

$$F(u, s) = F(u, \Lambda)F(s, \Lambda). \quad (18)$$

The form factor of Eq. (18) has to multiply the sum of both the  $s$  and  $u$  channel diagrams. Model D refers to calculations performed with the prescription Eq. (18).




FIG. 3: The data [10, 32] on the differential  $\gamma p \rightarrow \omega p$  cross section as a function of  $t$  for different photon energies  $E_\gamma$ . The lines show our calculations with different prescriptions for the  $\omega NN$  vertex functions given by model A (dashed), B (dotted), C (dashed-dotted), and D (solid).




FIG. 4: The differential  $\gamma p \rightarrow \omega p$  cross section as a function of  $t$  for  $E_\gamma = 2.8$  GeV. The data are from Refs. [10, 32]. The calculations were done by model B with parameters from Ref. [28] (a) and by model D (b). Dashed lines: contributions from the nucleon exchange; dotted lines: meson exchange; dashed-dotted lines: both nucleonic and meson exchanges.

Model D describes the  $d\sigma/dt$  and  $d\sigma/du$  data on  $\omega$  meson photoproduction at  $2.8 \leq E_\gamma \leq 4.7$  GeV better than all the other models (see e.g. the  $\chi^2$ -values) with the same number of parameters as is employed in model B. Moreover, we consider the parameters of model D as quite reasonable. However, our calculations could not reproduce the dip of the  $\gamma p \rightarrow \omega p$  differential cross section around  $|u| \simeq 0.15$  GeV<sup>2</sup> at the photon energy  $E_\gamma = 4.7$  GeV. One might speculate whether the appearance of a dip in  $d\sigma/du$  needs additional experimental confirmation.

As is shown in Fig. 3, the four different models reproduce the differential  $\omega$  photoproduction cross section at small  $|t|$  at different photon energies quite well, but substantially differ at  $|t| \geq 1$  GeV<sup>2</sup>. Therefore we conclude that the prescription of the  $\omega NN$  vertex function as well as the model parameters substantially influence the results in the region of moderate and large  $|t|$ .

To illustrate our finding explicitly, we calculate the  $\gamma p \rightarrow \omega p$  differential cross section  $d\sigma/dt$  by model B with the set of parameters from Ref. [28] taken as

$$\begin{aligned} g_{\pi NN} &= 13.26, \Lambda_{\pi NN} = 0.6, g_{\omega\gamma\pi} = 1.823, \Lambda_{\omega\gamma\pi} = 0.9, \\ g_{\eta NN} &= 3.53, \Lambda_{\eta NN} = 1.0, g_{\omega\gamma\eta} = 0.416, \Lambda_{\omega\gamma\eta} = 0.9 \\ g_{\omega NN} &= 10.35, \Lambda_{\omega NN} = 0.5, \end{aligned} \quad (19)$$

where the cut-off parameters are given in GeV. Furthermore, in Ref. [28] the pomeron exchange was included

instead of  $\sigma$  meson exchange but it was shown that the contribution from the pomeron exchange is negligible at  $E_\gamma=2.8$  GeV. Therefore we also neglect the contribution from the  $\sigma$  meson exchange in reproducing the results from Ref. [28] by model B.

Fig. 4a) shows the differential  $\gamma p \rightarrow \omega p$  cross section as a function of  $t$  for  $E_\gamma=2.8$  GeV. The calculations are based on model B with parameters given by Eq. (19). The dashed-dotted line shows the total contribution from both meson and nucleon exchanges, while the dashed line indicates the contribution from nucleon exchange only and the dotted one the contribution from mesonic exchanges. The calculations shown in Fig. 4a) are identical to the ones reported in Ref. [28]. It is clear that model B cannot reproduce the data at  $|t| \geq 0.5$  GeV<sup>2</sup>.




FIG. 5: The dependence of the differential  $\gamma p \rightarrow \omega p$  cross section  $d\sigma/du$  on the photon energy  $E_\gamma$ . The data were taken from Ref. [10]. The solid line shows our calculations based on model D.

Now, Fig. 4b) shows that the calculations based on model D reasonably reproduce the data [10] for the whole range of  $t$ -values. Comparing the two calculations one can easily notice that the contributions from mesonic currents from model B and model C are almost identical, whereas the contributions from the nucleon exchange are substantially different. Obviously, the prescription of the  $\omega NN$  vertex is important in understanding the data on  $\omega$  meson photoproduction at large and moderate  $|t|$ .

#### IV. ENERGY DEPENDENCE OF THE BACKWARD $\omega$ PHOTOPRODUCTION

Now we investigate the dependence of the differential  $\gamma p \rightarrow \omega p$  cross section  $d\sigma/du$  on the photon energy. Let us recall that the very steep  $s^{-8}$  dependence [11] was interpreted [10] as an indication of hard interactions between the photon and the quarks of the nucleon.

Experimental data [10] on the energy dependence of the  $\omega$  photoproduction cross section  $d\sigma/du$  are shown in Figs. 5-7. The solid lines show our calculations based on model D. The overall agreement between the data and the model calculations is reasonable, although there is a systematical discrepancy at  $-0.2 < u < -0.1$  GeV<sup>2</sup> for photon energies above 4 GeV, i.e. in the vicinity of the experimentally observed dip. Furthermore, the calculations systematically underestimate the part of the data at  $-0.8 < u < -0.4$  GeV<sup>2</sup>, but rather well reproduce the energy dependence of the differential  $\gamma p \rightarrow \omega p$  cross section  $d\sigma/du$ .

Two extreme situations are shown in Fig. 7. In Ref. [10] the  $\gamma p \rightarrow \omega p$  data at  $-0.1 < u < 0$  GeV<sup>2</sup> were fitted by a  $s^{-3.9 \pm 0.3}$  dependence, while experimental results at  $-1.8 < u < -1.6$  GeV<sup>2</sup> were fitted by a  $s^{-9.3 \pm 2.1}$  dependence. Other data available for the range  $-1.6 < u < -0.1$  GeV<sup>2</sup> indicate that the power of  $s$  is between the two extreme values given above.

Furthermore, our calculations can be fitted well by  $d\sigma/du \propto s^{4.57}$  at  $-0.1 < u < 0$  GeV<sup>2</sup> and  $d\sigma/du \propto s^{6.73}$  at  $-1.8 < u < -1.6$  GeV<sup>2</sup> respectively. Although the model does not produce the very steep  $s$  dependence, we conclude that the agreement between the calculations and the experimental data is very reasonable.

#### V. SUMMARY

The  $\omega$  meson photoproduction for energies  $2.8 \leq E_\gamma \leq 4.7$  GeV was studied within the boson exchange model. We consider the  $\pi$ ,  $\eta$ ,  $\sigma$  and nucleon exchanges. The mesonic exchanges dominate at small four momenta transfer squared  $|t|$ , while the nucleon exchange contributes at large  $|t|$ .

It was found that the  $\gamma p \rightarrow \omega p$  data [10] available at large  $|t|$  or small  $|u|$  are very sensitive to the prescription of the  $\omega NN$  vertex function. We investigated different recipes commonly used for restoration of gauge invariance in  $\omega$  photoproduction. The inclusion of a phenomenological form factor at the  $\omega NN$  vertex which is consistent with gauge invariance and crossing symmetry [38] allows to reproduce the data [10] both at small  $t$  and small  $u$ .

However we found that within our approach it is impossible to reproduce the dip in the differential  $\gamma p \rightarrow \omega p$  cross section  $d\sigma/du$  around  $u \simeq -0.15$  GeV<sup>2</sup>. Furthermore, since a similar dip was not observed in the backward photoproduction of  $\pi$ ,  $\rho$  and  $f$  mesons, we suggest that an additional experimental confirmation would be highly welcome.




FIG. 6: The dependence of the differential  $\gamma p \rightarrow \omega p$  cross section  $d\sigma/du$  on the photon energy  $E_\gamma$ . The data were taken from Ref. [10]. The solid line shows our calculations based on model D.

We examined the energy dependence of the differential  $\omega$  photoproduction cross section  $d\sigma/du$ . The experimentally observed steep  $\propto s^{-8}$  dependence was interpreted [10] as an indication [11] of hard interactions between the photon and quarks of the nucleon. However, our meson-exchange calculations also quite reasonably reproduce the energy dependence of  $d\sigma/du$ .

### Acknowledgments

Work of K.T. was supported by the Australian Research Council, and the Forschungszentrum Jülich. S.K. acknowledges support by grant N. 447AUS113/14/0 by the Deutsche Forschungsgemeinschaft. Thanks are given to Profs. J. Speth and A.W. Thomas for their support of this work.

- 
- [1] S.M. Berman and S.D. Drell, Phys. Rev. **133**, 3791 (1964).
  - [2] A. Donnachie and P.V. Landshoff, Phys. Lett. B **348**, 213 (1995).
  - [3] A. Donnachie and P.V. Landshoff, Phys. Lett. B **478**, 146 (2000).
  - [4] J.M. Laget, Phys. Lett. B **489**, 313 (2000).
  - [5] J. Breitweg et al., Eur. Phys. J. C **1**, 81 (1998).
  - [6] A. Donnachie and P.V. Landshoff, Phys. Lett. B **470**, 243 (1999).
  - [7] C. Adloff et al., Phys. Lett. B **483**, 23 (2000).
  - [8] A. Sibirtsev, K. Tsushima and A.W. Thomas, Phys. Rev. C **63**, 044906 (2001).
  - [9] A. Donnachie and P.V. Landshoff, Phys. Lett. B **518**, 63 (2001).
  - [10] R.W. Clift et al., Phys. Lett. B **72**, 144 (1977).
  - [11] S.J. Brodsky and G.R. Farrar, Phys. Rev. D **11**, 1309 (1975).
  - [12] R.W. Clift et al., Phys. Lett. B **64**, 213 (1976).
  - [13] R.L. Anderson et al., Phys. Rev. Lett. **23**, 721 (1969).
  - [14] D. Tompkins et al., Phys. Rev. Lett. **23**, 725 (1969).
  - [15] R. Machleidt, K. Holinde and C. Elster, Phys. Rept. **149**, 1 (1987).
  - [16] R. Machleidt, Phys. Rev. C **63**, 024001 (2001).
  - [17] S.D. Drell and F. Zachariasen, Phys. Rev. Lett. **5**, 66 (1960).
  - [18] W. Alles and D. Boccaletti, Nuov. Cim. **27**, 306 (1963).
  - [19] H. Joos and G. Kramer, Z. Phys. **178**, 542 (1964).
  - [20] K. Gottfried and J.D. Jackson, Nuov. Cim. **34**, 735 (1964).
  - [21] K.M. Watson, Phys. Lett. B **17**, 72 (1965).
  - [22] Aachen-Berlin-Bonn-Hamburg-Heidelberg-München Collaboration, Phys. Rev. **175**, 1669 (1969).
  - [23] J. Ballam et al., Proc. 4 Int. Symp. on Electron and Photon Interactions at High Energies, Daresbury (1969).
  - [24] J. Ballam et al., Phys. Rev. Lett. **24**, 1364 (1970).
  - [25] M. Davier et al., Phys. Rev. D **1**, 790 (1970).
  - [26] Landol-Börnstein, New Series **I/12b**.
  - [27] B. Friman and M. Soyeur, Nucl. Phys. A **600**, 477 (1996).
  - [28] Y. Oh, A.I. Titov and T.S.H. Lee, Phys. Rev. C **63**, 025201 (2001).
  - [29] A. Donnachie and P.V. Landshoff, Phys. Lett. B **185**, 403 (1987).
  - [30] A. Donnachie and P.V. Landshoff, Phys. Lett. B **296**, 227 (1992).
  - [31] F.J. Klein et al.,  $\pi N$  Newslett. **14**, 141 (1998).
  - [32] J. Ballam et al., Phys. Rev. D **7**, 3150 (1973).
  - [33] Particle Data Group, Eur. Phys. J. C **3**, 1 (1998).
  - [34] K. Ohta, Phys. Rev. C **40**, 1335 (1989).
  - [35] R.L. Workman, H.W.L. Naus and S.J. Pollock, Phys.




FIG. 7: The dependence of the differential  $\gamma p \rightarrow \omega p$  cross section  $d\sigma/du$  on the photon energy  $E_\gamma$ . The data were taken from Ref. [10]. The solid line shows our calculations based on model D.

- Rev. **45**, 2511 (1992).  
 [36] H. Haberzettl, Phys. Rev. C **56**, 2041 (1997)  
 [37] H. Haberzettl, C. Bennhold, T. Mart and T. Feuster, Phys. Rev. C **58**, 40 (1998)  
 [38] R.M. Davidson and R. Workman, Phys. Rev. C **63**, 025210 (2001).  
 [39] F. James and M. Roos, Comp. Phys. Comm. **10**, 343 (1975).